

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
25 January 2001 (25.01.2001)

PCT

(10) International Publication Number
WO 01/05421 A1

(51) International Patent Classification⁷: **A61K 38/16**,
39/02, 48/00, 45/06, 31/713, A61P 35/00

Molecular Microbiology, Calle Velazquez, 144, E-28006
Madrid (ES).

(21) International Application Number: PCT/GB00/02743

(74) Agents: **WALTON, Seán, M.** et al.; Mewburn Ellis, York
House, 23 Kingsway, London WC2B 6HP (GB).

(22) International Filing Date: 17 July 2000 (17.07.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
9916810.6 16 July 1999 (16.07.1999) GB

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(71) Applicant (*for all designated States except US*): **CANCER
RESEARCH CAMPAIGN TECHNOLOGY LIMITED**
[GB/GB]; Cambridge House, 6-10 Cambridge Terrace, Re-
gent's Park, London NW1 4JL (GB).

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **DE LA CUEVA
MENDEZ, Guillermo** [ES/ES]; Campus Universidad
Autonoma, Centro Nacional de Biotecnologia, Depart-
ment of Microbial Biotechnology, Cantoblanco, E-28049
Madrid (ES). **LASKEY, Ronald, Alfred** [GB/GB];
Wellcome/CRC Institute, Tennis Court Road, Cambridge
CB2 1QR (GB). **MILLS, Anthony, David** [GB/GB];
Wellcome/CRC Institute, Tennis Court Road, Cambridge
CB2 1QR (GB). **DIAZ OREJAS, Ramon** [ES/ES]; Cen-
tro de Investigaciones Biologicas-CSIG, Department of

Published:

- With international search report.
- Before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments.

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*



WO 01/05421 A1

(54) Title: METHODS EMPLOYING BACTERIAL TOXIN-ANTITOXIN SYSTEMS FOR KILLING EUKARYOTIC CELLS

(57) Abstract: A method of inhibiting cell proliferation and/or cell cycle progression, the method comprising providing within eukaryotic cells a bacterial toxin, in particular a bacterial toxin of a post-segregational killing system such as *ParD*. Activity or production of toxin and/or inhibitor, e.g. antidote, are controlled for selective killing of target cells, not-target cells. Target cells include tumour cells.

METHODS EMPLOYING BACTERIAL TOXIN-ANTITOXIN SYSTEMS FOR KILLING EUKARYOTIC CELLS

The present invention relates to killing cells, or at least
5 impeding cell cycle progression. More particularly it relates
to methods and means for attacking eukaryotic cells, such as
tumour cells, with cytostatic, cytotoxic and/or cytopathic
agents. Specifically, the present invention employs toxins
and toxin/antidote systems based on bacterial systems and
10 under appropriate regulation for inhibiting cellular growth
and preferably killing cells. In particular embodiments of
the invention killing is selective or specific for certain
target cells.

15 There are various contexts in which it is desired to kill
cells, in particular selectively to kill certain cells within
a population of cells. In some contexts inhibition of
cellular growth or proliferation, for instance by impeding
cell cycle progression, may be sufficient. For simplicity
20 herein, unless context provides otherwise, reference to
killing cells may be used to encompass such inhibition.

An important area of application is in treatment of tumours,
cancer, psoriasis, arteriosclerosis and other
25 hyper-proliferative disorders. Other applications of
embodiments of the present invention include targeting any
desired eukaryotic cell for killing or at least inhibition of
growth. This may include cell lineage knock-outs and targeted

cell ablation, for instance in developmental control, or organogenesis studies. *In vitro* applications include study of the control or replication in prokaryotic and/or eukaryotic cells, screening for an antidote for a toxin, or toxin
5 inhibited by an antidote, design of or screening for improved toxin and/or antidote factors, and analysis of physiological responses of different cell types to inhibition of cell progression and/or inhibition of DNA replication.

10 In plants, pathogen defence responses involve cell necrosis, for instance triggered at a site of pathogen infection or ingress. Induced resistance is strongly correlated with the hypersensitive response (HR), an induced response associated with localized cell death at sites of attempted pathogen
15 ingress. It is hypothesized that by HR the plant deprives the pathogen of living host cells.

Many plant defence mechanisms are strongly induced in response to a challenge by an unsuccessful pathogen. Such an induction
20 of enhanced resistance can be systemic. It is believed that when a plant is challenged by a pathogen to which it is resistant, it undergoes an HR at the site of attempted ingress of the incompatible pathogen. The induced HR leads to a systemic enhancement and acquisition of plant resistance to
25 virulent pathogens that would normally cause disease in the unchallenged plant.

Artificial induction of cell death in plants has been shown to

be able to provide pathogen resistance, even where the mechanism inducing cell death is not triggered by any pathogen resistance gene. For instance, genes coding for substances leading to rapid cell death, such as BARNASE or diphtheria toxin may be use to induce the changes that lead to acquired resistance even though cell death in these latter examples is not caused by activation of the defence response. BARNASE is a ribonuclease from *Bacillus amyloliquifaciens* (Hartley (1988) J. Mol. Biol. 202: 913-915; Hartley (1989) Trends Biochem. Sci. 14: 450-454) and there is a corresponding protein called BARSTAR which inhibits BARNASE by forming a complex with it.

Use of embodiments of the present invention in plants may be used to generate protection against attack from fungi, bacteria, viruses or nematodes.

Plants of particular interest for use in embodiments of the present include cereals, maize, corn, wheat, barley, oats, rice, Brassicas, curcubits, potatoes, tomatoes, cotton, soya bean, and carrot.

Another use of embodiments of the present invention in plants include generation of male sterility (Mariani et al. Nature 357 384-387). For instance toxin or a toxin system in accordance with the present invention may be introduced into plants under appropriate control for tapetal-specific expression (Seurinck et al. (1990) Nucleic Acids Res. 18: 3403; Koltunow et al. (1990) Plant Cell 2, 1201-1224; Mariani et al (1990) Nature

347: 737-741). Male sterility in plants facilitates hybrid seed generation by preventing self-pollination, allowing agriculturalists to take advantage of so-called "hybrid vigour" by which crosses between inbred plant lines often
5 result in progeny with higher yield and increased resistance to disease. Provision of horticultural or ornamental plants lacking ability to make pollen may be used to reduce allergy problems of local inhabitants or for aesthetic reasons (e.g. in lilies, where anthers are currently removed by hand).

10

A further use in plants is in generation of seedlessness, often desirable for convenience and taste in produce such as watermelons, grapes, oranges and related fruits, tomatoes, peppers, cucumbers and so on. Toxin can be placed under
15 regulatory control of a seed-specific promoter, such as the promoter of a seed storage protein (Higgins et al, (1984) *Ann. Rev. Plant. Physiol.* 35: 191-221; Goldberg et al (1989) *Cell* 56: 149-160). Examples of seed-specific promoters include those for bean β -phaseolin (Sengupta-Gopalan et al, (1985)
20 *PNAS USA* 82: 3320-3324), bean lectin (Voelker et al (1987) *EMBO J.* 6: 3571-3577), soybean lectin (Ocamuro et al. (1986) *PNAS USA* 83: 8240-8344), rapeseed napin (Radke et al. *Theor. Appl. Genet.* 75: 685-694), maize zein (Hoffman et al (1987) *EMBO J.* 6: 3213-3221), barley β -hordein (Marris et al (1988) *Plant*
25 *Mol. Biol.* 10: 359-366) and wheat glutenin (Colot et al. (1987) *EMBO J.* 6: 3559-3564).

Prokaryotic plasmids have developed different genetic systems

that increase their stable maintenance in bacterial hosts. These systems are classified into two different types: partition systems, that ensure a well controlled partition of plasmid DNA copies between the two daughter cells, and killer systems, that eliminate from the bacterial population those daughter cells that have lost the plasmid during division (Yarmolinsky, Science (1995) Feb 10, 267(5199): 836-7). The latter are composed of two components: a bacterial toxin (always a protein), and its antidote (a protein or an antisense RNA that inhibits transcription of its killer partner) (Jensen and Gerdes, Mol. Microbiol. (1995) Jul. 17(2): 205-10; Thisted et al. J. Mol. Biol. (1992). Jan. 5 223(1):41-54). These killer systems are generally organized similarly from a molecular point of view, and several mechanisms ensure that a typical killer system is not activated if the stability of its harbouring plasmid is not compromised. Thus, both proteic antidote and toxic components are organized in a bicistronic operon, and the system is molecularly designed in such a way that both transcriptional and translational processes are optimised to maintain it in a silent state (i.e. a state in which the toxic component is being neutralised by its antidote) (Jensen and Gerdes, Mol. Microbiol. (1995) Jul 17(2): 205-10; Holcik and Iyer, Microbiology (1997), 143: 3403-3416).

25

Under normal circumstances, both components of a killer system are synthesized at a basal level in the host by its harbouring plasmid, allowing the host to survive. If a segregant

bacteria (i.e. a bacteria that has lost the plasmid) appears after cell division, another characteristic of these systems allows activation of the killing process in order to counter-select that specific cell: that is, the stability of the
5 antidote is lower than the toxin. Thus, without a continuous synthesis of the antidote, its preferential degradation leads to the appearance of a non-neutralised toxin that is then able to exert its lethal effect over the host. This toxic effect can be executed affecting different cellular targets,
10 depending on the specific killer system, for example DnaB dependent replication (parD, pem), DNA-gyrase complex (ccd), protein synthesis inhibition (KicB), and septum formation (kil), (for references see Holcik and Iyer, Microbiology (1997), 143, 3403-3416). Yarmolinsky describes in Science,
15 Vol. 267 (1995) other putative "addiction molecules" like the type II restriction enzymes (putative toxins) Pae R7 and EcoRI and their cognate methylases, that enhance the apparent stability of their harbouring plasmids (the original reference for this addiction modules is in Naito et al. Science 267:897
20 (1995)). In this work, Yarmolinsky also describes a couple of putative killer systems from bacteriophage lambda (Rex protein) and a couple of strains of *E. coli* carrying the gene cluster prr, that encodes for an anticodon nuclease that can be activated by a 26 residue polypeptide from bacteriophage T4
25 and can then cleave a transfer RNA important for lysine incorporation into proteins. T4 is invulnerable to this protein because it encodes for a couple of otherwise non-essential proteins that undoes the damage. He also describes

strains of *E. coli* that carry defective prophage ϕ 14, and that accomplish exclusion by cleavage of elongation factor Tu and inhibiting translation globally.

5 ParD is one of these killer systems (Bravo et al. Mol. Gen. Genet. (1987) Nov. 210(1): 101-10; Bravo et al. Mol. Gen. Genet. (1988). Dec. 215(1): 146-51). It is encoded by Gram negative plasmid R1 and is composed of two genes: *kis* (for killing suppressor) and *kid* (for killing determinant) that
10 encode for the antidote (10 KDa) and the toxin (12 KDa) respectively. ParD is a cryptic killer system that is tightly regulated to avoid its activation under circumstances that do not compromise R1 stability. Thus, it is controlled by coupled transcription (Ruiz-Echevarria et al. Mol. Microbiol.
15 (1991) Nov. 5(11): 2685-93), by post-transcriptional processing of its bicistronic mRNA (Ruiz-Echevarria et al. Mol. Gen. Genet. (1995) Sep. 20 248(5): 599-609), by overlapped translation (Ruiz-Echevarria et al. Mol. Gen. Genet. (1995) Sep. 20 248(5): 599-609), and by a very tight
20 interaction between *Kis* and *Kid* to form a non-toxic complex that, at the same time, is able to repress transcription from its own promoter (Ruiz-Echevarria et al. Mol. Microbiol. (1991) Nov. 5(11): 2685-93). Genetic organisation of *ParD* favours coupled transcription, overlapped translation and
25 post-transcriptional modification of some of the obtained mRNA. *Kis/Kid* complexes repress transcription of *kis* and *kid* genes.

ParD homologues have been described at least in plasmid R100 (pem system) (Tsuchimoto et al. J. Bacteriol. (1988) Apr. 170(4): 1461-6; Tsuchimoto et al. J. Bacteriol. (1992) Jul. 174(13): 4205-11; Tsuchimoto et al. Mol. Gen. Genet. (1993) 5 Feb. 237(1-2): 81-88); Masuda et al. J. Bacteriol. (1993) Nov. 175(21): 6850-6) and in *E. coli* chromosome (ChpA and ChpB systems) (Tsuchimoto et al. Mol. Gen. Genet. (1993) Feb. 237(1-2): 81-88). Others are revealed by database searching.

10 Kid inhibits initiation of replication of the *E. coli* genome and of DnaB (i.e. the main replicative helicase of *E. coli*) dependent replication plasmids (Ruiz-Echevarria et al. J. Mol. Biol. (1995) Apr. 7 247(4): 568-77), and over-expression of the latter titrates the toxic effect of the former in this
15 organism *in vivo* (Ruiz-Echevarria et al. J. Mol. Biol. (1995) Apr. 7 247(4): 568-77), suggesting that DnaB is involved in the mechanism of inhibition by Kid. Recent observations in the inventors' laboratory strongly suggest that this inhibition is due neither to disassembly by Kid of DnaB
20 hexameric complexes in solution nor to inhibition of its helicase activity over a wide range of substrates including *oriC*, the replication origin of the *E. coli* genome. Without wishing to be limited by theory, it may be that loading of DnaB at the origin of replication is the process inhibited by
25 Kid, either by direct interaction between them and/or mediated by a third component (DNA or protein) yet to be described. Current research is focused on the identification of the exact mechanism of action of Kid from a molecular point of view.

Until the work of the present inventors disclosed herein it was not obvious that prokaryotic systems that have evolved for specific roles in bacteria could function in eukaryotic cells.

5 For instance, in a two-component killer system such as involving *kis/kid*, both components need to perform their respective functions - the toxin to kill cells in the absence of antidote (or when present in excess of antidote), and the antidote to both neutralise the toxin and be controllable, for
10 instance by a mechanism involving rapid turnover. Preferably the toxin does not exert any side effect on cell viability. Rather, it is preferred that cell killing is via a programmed cell death mechanism such as apoptosis. In plants it may be preferred for certain applications to induce a necrotic
15 response, e.g. in inducing or enhancing pathogen resistance.

The present inventors have shown that bacterial toxin and antidote are functional in eukaryotic cells, yeast, *Xenopus* and mammalian (in particular human), and can be controlled to
20 inhibit cell cycle progression and cellular proliferation and to kill cells. It is shown in experiments described below that cells can be killed by apoptosis.

Brief Description of the Figures

25 Figure 1 shows results of experiments showing that a promoter induced by Cu^{2+} later used for control of *Kis* antidote expression and a different promoter repressed by methionine later used for control of *Kid* toxin expression are both

functional in *S. cerevisiae*. The graph shows units of β -galactosidase activity at different concentrations (μ M) of the regulatory factors Cu^{2+} (light circles) and Met (dark circles).

- 5 Figure 2 shows results of experiments demonstrating the effect of Doxycyclin on a Tetracyclin regulatable promoter (Tet Pr) activity in HeLa cells (light bars), this promoter later used for control of expression of antidote Kis (in the vector pTRE-Luc), and an absence of effect of Doxycyclin on
- 10 Cytomegalovirus Early promoter (CMV Pr) activity (dark bars), this promoter later used for control of expression of toxin Kid (pCMV-Luc). Luciferase activity is plotted, in arbitrary units.
- 15 Figure 3 illustrates various constructs employed for expression of Kis and/or Kid in HeLa cells.

Figure 4 shows results of experiments in which *kis* expression was modulated by Doxycyclin in cultures of HeLa cells stably

20 transfected with pNATHA1i and pNATHA2i (Figure 3). Kid expression was controlled by CMV Pr which is unaffected by Doxycyclin.

Figure 5 shows further results of experiments (numbers of dead

25 cells) in which *kis* expression was modulated by Doxycyclin in cultures of HeLa cells stably transfected with pNATHA1i and pNATHA2i (Figure 5). Kis expression was controlled by the Tet Pr which is repressed by Doxycyclin, while *kid* expression was

controlled by CMV Pr which is unaffected by Doxycyclin.

Figure 6 shows emergence of the apoptosis marker Annexin V in cells subject to the experiments of which results are shown in Figure 5, indicating the cell death caused by Kid to involve apoptosis.

According to one aspect of the present invention there is provided a method of inhibiting cell proliferation and/or cell cycle progression, the method comprising providing within eukaryotic cells a bacterial toxin. The bacterial toxin is generally a toxin of a bacterial cell killing system, preferably of a post-segregational killing system. As is explained herein, these are mostly plasmid-borne in bacteria although some are found on the bacterial chromosome, and others functional in bacterial cells are encoded by bacteriophage.

Preferably a toxin of use in the present invention interferes with DNA replication, and thereby impedes cell cycle progression and/or triggers programmed cell death. As noted, other physiological processes may be inhibited by means of other toxins. The target of the toxin may be *DnaB* or DNA gyrase.

25

Preferably a bacterial toxin employed in the present invention triggers programmed cell death. Experiments described below

demonstrate use of bacterial toxin to induce apoptosis in mammalian cells.

Some measure of control of toxin action is preferably employed in aspects of the present invention. Bacterial cell killing systems of use in the present invention naturally employ an antidote to the toxin. The present inventors have shown that both toxin and antidote of a bacterial cell killing system are functional in various eukaryotic cells and that their
5
10
15
20
respective activities can be controlled for selective inhibition of cellular proliferation or impedance of cell cycle progression, and/or induction of programmed cell death.

A bacterial cell killing system employed in the present invention may comprise a toxin and an antidote which are both protein. Such a killing system is termed in the art a "proteic killer gene system" - Jensen & Gerdes, 1995, Mol. Microbiol. (1995) Jul 17(2): 205-10). A bacterial killer system of use in the invention may be an *E. coli* system or
20
other bacterium.

Examples of bacterial killer systems of use in the present invention, and comprising toxins of use in the present invention (for references see Holcik and Iyer (1997),
25
30
35
40
45
50
55
60
65
70
75
80
85
90
95
100
105
110
115
120
125
130
135
140
145
150
155
160
165
170
175
180
185
190
195
200
205
210
215
220
225
230
235
240
245
250
255
260
265
270
275
280
285
290
295
300
305
310
315
320
325
330
335
340
345
350
355
360
365
370
375
380
385
390
395
400
405
410
415
420
425
430
435
440
445
450
455
460
465
470
475
480
485
490
495
500
505
510
515
520
525
530
535
540
545
550
555
560
565
570
575
580
585
590
595
600
605
610
615
620
625
630
635
640
645
650
655
660
665
670
675
680
685
690
695
700
705
710
715
720
725
730
735
740
745
750
755
760
765
770
775
780
785
790
795
800
805
810
815
820
825
830
835
840
845
850
855
860
865
870
875
880
885
890
895
900
905
910
915
920
925
930
935
940
945
950
955
960
965
970
975
980
985
990
995
1000
1005
1010
1015
1020
1025
1030
1035
1040
1045
1050
1055
1060
1065
1070
1075
1080
1085
1090
1095
1100
1105
1110
1115
1120
1125
1130
1135
1140
1145
1150
1155
1160
1165
1170
1175
1180
1185
1190
1195
1200
1205
1210
1215
1220
1225
1230
1235
1240
1245
1250
1255
1260
1265
1270
1275
1280
1285
1290
1295
1300
1305
1310
1315
1320
1325
1330
1335
1340
1345
1350
1355
1360
1365
1370
1375
1380
1385
1390
1395
1400
1405
1410
1415
1420
1425
1430
1435
1440
1445
1450
1455
1460
1465
1470
1475
1480
1485
1490
1495
1500
1505
1510
1515
1520
1525
1530
1535
1540
1545
1550
1555
1560
1565
1570
1575
1580
1585
1590
1595
1600
1605
1610
1615
1620
1625
1630
1635
1640
1645
1650
1655
1660
1665
1670
1675
1680
1685
1690
1695
1700
1705
1710
1715
1720
1725
1730
1735
1740
1745
1750
1755
1760
1765
1770
1775
1780
1785
1790
1795
1800
1805
1810
1815
1820
1825
1830
1835
1840
1845
1850
1855
1860
1865
1870
1875
1880
1885
1890
1895
1900
1905
1910
1915
1920
1925
1930
1935
1940
1945
1950
1955
1960
1965
1970
1975
1980
1985
1990
1995
2000
2005
2010
2015
2020
2025
2030
2035
2040
2045
2050
2055
2060
2065
2070
2075
2080
2085
2090
2095
2100
2105
2110
2115
2120
2125
2130
2135
2140
2145
2150
2155
2160
2165
2170
2175
2180
2185
2190
2195
2200
2205
2210
2215
2220
2225
2230
2235
2240
2245
2250
2255
2260
2265
2270
2275
2280
2285
2290
2295
2300
2305
2310
2315
2320
2325
2330
2335
2340
2345
2350
2355
2360
2365
2370
2375
2380
2385
2390
2395
2400
2405
2410
2415
2420
2425
2430
2435
2440
2445
2450
2455
2460
2465
2470
2475
2480
2485
2490
2495
2500
2505
2510
2515
2520
2525
2530
2535
2540
2545
2550
2555
2560
2565
2570
2575
2580
2585
2590
2595
2600
2605
2610
2615
2620
2625
2630
2635
2640
2645
2650
2655
2660
2665
2670
2675
2680
2685
2690
2695
2700
2705
2710
2715
2720
2725
2730
2735
2740
2745
2750
2755
2760
2765
2770
2775
2780
2785
2790
2795
2800
2805
2810
2815
2820
2825
2830
2835
2840
2845
2850
2855
2860
2865
2870
2875
2880
2885
2890
2895
2900
2905
2910
2915
2920
2925
2930
2935
2940
2945
2950
2955
2960
2965
2970
2975
2980
2985
2990
2995
3000
3005
3010
3015
3020
3025
3030
3035
3040
3045
3050
3055
3060
3065
3070
3075
3080
3085
3090
3095
3100
3105
3110
3115
3120
3125
3130
3135
3140
3145
3150
3155
3160
3165
3170
3175
3180
3185
3190
3195
3200
3205
3210
3215
3220
3225
3230
3235
3240
3245
3250
3255
3260
3265
3270
3275
3280
3285
3290
3295
3300
3305
3310
3315
3320
3325
3330
3335
3340
3345
3350
3355
3360
3365
3370
3375
3380
3385
3390
3395
3400
3405
3410
3415
3420
3425
3430
3435
3440
3445
3450
3455
3460
3465
3470
3475
3480
3485
3490
3495
3500
3505
3510
3515
3520
3525
3530
3535
3540
3545
3550
3555
3560
3565
3570
3575
3580
3585
3590
3595
3600
3605
3610
3615
3620
3625
3630
3635
3640
3645
3650
3655
3660
3665
3670
3675
3680
3685
3690
3695
3700
3705
3710
3715
3720
3725
3730
3735
3740
3745
3750
3755
3760
3765
3770
3775
3780
3785
3790
3795
3800
3805
3810
3815
3820
3825
3830
3835
3840
3845
3850
3855
3860
3865
3870
3875
3880
3885
3890
3895
3900
3905
3910
3915
3920
3925
3930
3935
3940
3945
3950
3955
3960
3965
3970
3975
3980
3985
3990
3995
4000
4005
4010
4015
4020
4025
4030
4035
4040
4045
4050
4055
4060
4065
4070
4075
4080
4085
4090
4095
4100
4105
4110
4115
4120
4125
4130
4135
4140
4145
4150
4155
4160
4165
4170
4175
4180
4185
4190
4195
4200
4205
4210
4215
4220
4225
4230
4235
4240
4245
4250
4255
4260
4265
4270
4275
4280
4285
4290
4295
4300
4305
4310
4315
4320
4325
4330
4335
4340
4345
4350
4355
4360
4365
4370
4375
4380
4385
4390
4395
4400
4405
4410
4415
4420
4425
4430
4435
4440
4445
4450
4455
4460
4465
4470
4475
4480
4485
4490
4495
4500
4505
4510
4515
4520
4525
4530
4535
4540
4545
4550
4555
4560
4565
4570
4575
4580
4585
4590
4595
4600
4605
4610
4615
4620
4625
4630
4635
4640
4645
4650
4655
4660
4665
4670
4675
4680
4685
4690
4695
4700
4705
4710
4715
4720
4725
4730
4735
4740
4745
4750
4755
4760
4765
4770
4775
4780
4785
4790
4795
4800
4805
4810
4815
4820
4825
4830
4835
4840
4845
4850
4855
4860
4865
4870
4875
4880
4885
4890
4895
4900
4905
4910
4915
4920
4925
4930
4935
4940
4945
4950
4955
4960
4965
4970
4975
4980
4985
4990
4995
5000
5005
5010
5015
5020
5025
5030
5035
5040
5045
5050
5055
5060
5065
5070
5075
5080
5085
5090
5095
5100
5105
5110
5115
5120
5125
5130
5135
5140
5145
5150
5155
5160
5165
5170
5175
5180
5185
5190
5195
5200
5205
5210
5215
5220
5225
5230
5235
5240
5245
5250
5255
5260
5265
5270
5275
5280
5285
5290
5295
5300
5305
5310
5315
5320
5325
5330
5335
5340
5345
5350
5355
5360
5365
5370
5375
5380
5385
5390
5395
5400
5405
5410
5415
5420
5425
5430
5435
5440
5445
5450
5455
5460
5465
5470
5475
5480
5485
5490
5495
5500
5505
5510
5515
5520
5525
5530
5535
5540
5545
5550
5555
5560
5565
5570
5575
5580
5585
5590
5595
5600
5605
5610
5615
5620
5625
5630
5635
5640
5645
5650
5655
5660
5665
5670
5675
5680
5685
5690
5695
5700
5705
5710
5715
5720
5725
5730
5735
5740
5745
5750
5755
5760
5765
5770
5775
5780
5785
5790
5795
5800
5805
5810
5815
5820
5825
5830
5835
5840
5845
5850
5855
5860
5865
5870
5875
5880
5885
5890
5895
5900
5905
5910
5915
5920
5925
5930
5935
5940
5945
5950
5955
5960
5965
5970
5975
5980
5985
5990
5995
6000
6005
6010
6015
6020
6025
6030
6035
6040
6045
6050
6055
6060
6065
6070
6075
6080
6085
6090
6095
6100
6105
6110
6115
6120
6125
6130
6135
6140
6145
6150
6155
6160
6165
6170
6175
6180
6185
6190
6195
6200
6205
6210
6215
6220
6225
6230
6235
6240
6245
6250
6255
6260
6265
6270
6275
6280
6285
6290
6295
6300
6305
6310
6315
6320
6325
6330
6335
6340
6345
6350
6355
6360
6365
6370
6375
6380
6385
6390
6395
6400
6405
6410
6415
6420
6425
6430
6435
6440
6445
6450
6455
6460
6465
6470
6475
6480
6485
6490
6495
6500
6505
6510
6515
6520
6525
6530
6535
6540
6545
6550
6555
6560
6565
6570
6575
6580
6585
6590
6595
6600
6605
6610
6615
6620
6625
6630
6635
6640
6645
6650
6655
6660
6665
6670
6675
6680
6685
6690
6695
6700
6705
6710
6715
6720
6725
6730
6735
6740
6745
6750
6755
6760
6765
6770
6775
6780
6785
6790
6795
6800
6805
6810
6815
6820
6825
6830
6835
6840
6845
6850
6855
6860
6865
6870
6875
6880
6885
6890
6895
6900
6905
6910
6915
6920
6925
6930
6935
6940
6945
6950
6955
6960
6965
6970
6975
6980
6985
6990
6995
7000
7005
7010
7015
7020
7025
7030
7035
7040
7045
7050
7055
7060
7065
7070
7075
7080
7085
7090
7095
7100
7105
7110
7115
7120
7125
7130
7135
7140
7145
7150
7155
7160
7165
7170
7175
7180
7185
7190
7195
7200
7205
7210
7215
7220
7225
7230
7235
7240
7245
7250
7255
7260
7265
7270
7275
7280
7285
7290
7295
7300
7305
7310
7315
7320
7325
7330
7335
7340
7345
7350
7355
7360
7365
7370
7375
7380
7385
7390
7395
7400
7405
7410
7415
7420
7425
7430
7435
7440
7445
7450
7455
7460
7465
7470
7475
7480
7485
7490
7495
7500
7505
7510
7515
7520
7525
7530
7535
7540
7545
7550
7555
7560
7565
7570
7575
7580
7585
7590
7595
7600
7605
7610
7615
7620
7625
7630
7635
7640
7645
7650
7655
7660
7665
7670
7675
7680
7685
7690
7695
7700
7705
7710
7715
7720
7725
7730
7735
7740
7745
7750
7755
7760
7765
7770
7775
7780
7785
7790
7795
7800
7805
7810
7815
7820
7825
7830
7835
7840
7845
7850
7855
7860
7865
7870
7875
7880
7885
7890
7895
7900
7905
7910
7915
7920
7925
7930
7935
7940
7945
7950
7955
7960
7965
7970
7975
7980
7985
7990
7995
8000
8005
8010
8015
8020
8025
8030
8035
8040
8045
8050
8055
8060
8065
8070
8075
8080
8085
8090
8095
8100
8105
8110
8115
8120
8125
8130
8135
8140
8145
8150
8155
8160
8165
8170
8175
8180
8185
8190
8195
8200
8205
8210
8215
8220
8225
8230
8235
8240
8245
8250
8255
8260
8265
8270
8275
8280
8285
8290
8295
8300
8305
8310
8315
8320
8325
8330
8335
8340
8345
8350
8355
8360
8365
8370
8375
8380
8385
8390
8395
8400
8405
8410
8415
8420
8425
8430
8435
8440
8445
8450
8455
8460
8465
8470
8475
8480
8485
8490
8495
8500
8505
8510
8515
8520
8525
8530
8535
8540
8545
8550
8555
8560
8565
8570
8575
8580
8585
8590
8595
8600
8605
8610
8615
8620
8625
8630
8635
8640
8645
8650
8655
8660
8665
8670
8675
8680
8685
8690
8695
8700
8705
8710
8715
8720
8725
8730
8735
8740
8745
8750
8755
8760
8765
8770
8775
8780
8785
8790
8795
8800
8805
8810
8815
8820
8825
8830
8835
8840
8845
8850
8855
8860
8865
8870
8875
8880
8885
8890
8895
8900
8905
8910
8915
8920
8925
8930
8935
8940
8945
8950
8955
8960
8965
8970
8975
8980
8985
8990
8995
9000
9005
9010
9015
9020
9025
9030
9035
9040
9045
9050
9055
9060
9065
9070
9075
9080
9085
9090
9095
9100
9105
9110
9115
9120
9125
9130
9135
9140
9145
9150
9155
9160
9165
9170
9175
9180
9185
9190
9195
9200
9205
9210
9215
9220
9225
9230
9235
9240
9245
9250
9255
9260
9265
9270
9275
9280
9285
9290
9295
9300
9305
9310
9315
9320
9325
9330
9335
9340
9345
9350
9355
9360
9365
9370
9375
9380
9385
9390
9395
9400
9405
9410
9415
9420
9425
9430
9435
9440
9445
9450
9455
9460
9465
9470
9475
9480
9485
9490
9495
9500
9505
9510
9515
9520
9525
9530
9535
9540
9545
9550
9555
9560
9565
9570
9575
9580
9585
9590
9595
9600
9605
9610
9615
9620
9625
9630
9635
9640
9645
9650
9655
9660
9665
9670
9675
9680
9685
9690
9695
9700
9705
9710
9715
9720
9725
9730
9735
9740
9745
9750
9755
9760
9765
9770
9775
9780
9785
9790
9795
9800
9805
9810
9815
9820
9825
9830
9835
9840
9845
9850
9855
9860
9865
9870
9875
9880
9885
9890
9895
9900
9905
9910
9915
9920
9925
9930
9935
9940
9945
9950
9955
9960
9965
9970
9975
9980
9985
9990
9995
10000
10005
10010
10015
10020
10025
10030
10035
10040
10045
10050
10055
10060
10065
10070
10075

system such as *ParD* (of R1 or homologues as discussed above, *ccdA* (H or *let A*) of the F plasmid (antidote) and *ccdB* (G, *letB* or *letD*) toxin which acts by poisoning DNA-gyrase complexes (Jaffé, et al. (1985), *Bacteriol*, 163: 841-849) note
5 that the mode of action of the *ParD* system is remarkably similar to that of the *Ccd* system), bacteriophage P1 toxic protein *Doc* with antidote *Phd* (Lehnherr, et al. (1993), *Mol. Biol.*, 233: 414-428), *parDE* of plasmid RK2 (Roberts et al., 1994 *J. Mol. Biol.* 268, 27109-27117), with toxic protein *ParE*
10 and antidote *ParD*, and *hig* of plasmid Rts1 (Tian et al., 1996, *Biochem biophys Res Commun* 220 280-284) with antidote *higA* to toxin *higB*.

Further examples of bacterial killer systems of use in the
15 present invention and comprising toxins of use in the present invention, where the natural antidote is an antisense RNA include *parB* of plasmid R1 (Gerdes, et al. (1990a), *New Biol*, 2: 946-956) with toxin *Hok* and antidote *Sok* (Thisted et al, 1994, *EMBO J.* 13, 1950-1959; *hok* mRNA is very stable but *sok*
20 RNA decays rapidly), *srnB* (Onishi, (1975), *Science*, 187: 257-258) *flm* (Loh, et al. (1988), *Gene*, 66: 259-268) of the F plasmid and *pnd* of both IncI plasmid R483 and IncB plasmid R16 (Akimoto and Ohnishi (1982), *Microbiol. Immunol.*, 26: 779-793), *relF* of the *E. coli* chromosomal *relB* operon (induction
25 of the *relF* gene leads to the same physiological response as expression of the *hok* gene - Gerdes, et al. (1986a), *EMBO J.* 5: 2023-2029), *relB* homologues (Gronlund and Gerdes, 1999, *J. Mol. Biol.* 285, 1401-1415) and *Gef* (also chromosomal) which is

structurally and functionally similar to the proteins encoded by *hok* and *relF* (Poulsen, et al. (1989), *Mol. Microbiol.*, 3: 1463-1472). Gef protein is toxic and regulated by antisense RNA Sof.

5

Further systems of use and comprising toxins of use in the present invention include *SegB* operon epsilon (antidote) and zeta (toxin) of pSM19035 and pDB101 (Ceglowski et al. (1993) *Mol. Gen. Genet.* 241(5-6): 579-85; Ceglowski et al. (1993) 10 *Gene* 136(1-2): 1-12), *kicA* (antidote) and *kicB* (toxin) found in the *E. coli* chromosome (Feng, et al. (1984), *Mol. Gen. Genet.*, 243: 136-147), and the *kil/kor* systems carried by bacterial plasmids of the incompatibility groups P and N. See Holcik and Iyer (*Microbiology* (1997) 143: 3403-3416) for 15 examples and references. See also Jensen and Gerdes (*Mol. Microbiol.* (1995) 17(2), 205-210) and Yarmolinsky (*Science*, (1995) 267, 836-837) for reviews of proteic killer gene systems, noted to have striking similarities in both structure and function.

20

The toxin of any of these systems may be employed with the respective antidote. Alternatively or additionally, the toxin may be employed with one or more other elements which inhibit or block its activity (which may be by inhibiting or blocking 25 its production) as discussed.

In preferred embodiments both toxin and antidote of a bacterial cell killing system as disclosed, or toxin and other

inhibitor of its activity (e.g. inhibitor of its production) are introduced into eukaryotic cells under appropriate control for selective cell cycle inhibition and/or killing.

5 A method of the invention may include providing bacterial toxin and antidote or other toxin inhibitor to eukaryotic cells and, in target cells, removing or inhibiting the antidote or inhibitor to allow the toxin to work. Production or activity of antidote or inhibitor may be inhibited or
10 blocked. This may be by provision of an appropriate stimulus, e.g. inducer or repressor molecule of a promoter controlling antidote production, or may occur under conditions prevailing in target cells. As discussed below, the presence of a different form of a protein such as p53 in target cells vs.
15 non-target cells (e.g. for p53 tumour and non-tumour cells) can be employed as a controlling stimulus. An inducer or repressor molecule may be delivered to target cells to inhibit or block antidote and/or upregulate toxin.

20 Generally, the cell killing system is provided to cells by means of nucleic acid encoding the relevant components and, where applicable, control elements (discussed further below).

Control elements may include any one or more of those
25 available in the art allowing for selective variation of the ratio of toxin versus antidote. Examples include an inducible, repressible or constitutive promoter, antisense constructs and their activator or repressors, ribozymes,

splicing sequences and splicing factors, recombination systems (e.g. Cre-lox or FLP), wild-type or modified Internal Ribosome Entry Sites (IRES) (Schmid and Wimmer (1994), *Arch. Virol. Suppl.*, 9: 279-89; Borman, et al. (1994), *EMBO J.*, 1:13(13): 5 3149-57) and IRES inhibitors such as a yeast RNA that inhibits entry of ribosomes at some IRES (Das, et al. (1996), *J. Virol.*, 70(3): 1624-32; Das, et al. (1998), *J. Virol.*, 72(7): 6638-47; Das, et al. (1998), *Front Biosci.*, 1:3: D1241-52; Venkatosan, et al. (1999), *Nucleic Acids Res.* 15:27(2): 562- 10 72), elements that allow transcriptional interference between promoters (Greger and Proudfoot (1998), 17:17(16): 4771-9; Eggermont and Proudfoot (1993), *EMBO J.*, 12(6): 2539-48; Bateman and Paule (1998), *Cell*, 23:54(7): 985-92; Ponnambalam and Busby (1987), *FEBS Lett.*, 9:212(1): 21-7; Greger, et al. 15 (1998), *Nucleic Acids Res.*, 1:26(5): 1294-301), inteins (Chong et al. (1996) *J. Biol Chem* 271(16): 22159-68).

Activity of a bacterial toxin may be controlled by control of its production by expression from nucleic acid under control 20 of a regulatable promoter. It may be controlled by means of its natural antidote, which may be a protein or RNA. As noted, some natural antidotes are antisense RNAs that regulate production of toxin. Artificial antidotes or inhibitors may be designed and employed to control activity of any toxin. 25 So, for example, an antisense RNA or ribozyme may be designed to inhibit or block production of any toxin, even where the natural antidote of the toxin is a protein. A further option is to employ instead of natural antidote a different protein

that inhibits the toxin, for instance a protein (such as an antibody or binding fragment) that can be intracellularly expressed and which will bind the toxin within cells to neutralise its action. Any one or more of these various
5 approaches can be applied as alternatives or in combination.

One further aspect of the present invention provides a eukaryotic vector comprising nucleic acid encoding a toxin or cell killing system as disclosed. Such a vector may be used
10 to provide the toxin or cell killing system to eukaryotic cells.

Nucleic acid encoding a bacterial toxin and antidote may be provided as part of a vector or vectors suitable for
15 transformation of eukaryotic cells. Preferably the vector is suitable for transformation of target cells, for instance it may be suitable for transformation of plant cells (e.g. an *Agrobacterium* vector). Where two components of a bacterial killing system are employed, or a toxin is employed and a
20 specifically designed regulatory element is employed (e.g. antisense or ribozyme), preferably both components and regulatory elements for control of expression are provided on the same vector, but may be provided on separate vectors. Either or both of the encoding nucleotide sequences may be
25 under transcriptional control of a specific and/or regulatable promoter. Toxin- and antidote- encoding sequences may be provided in a "tail-to-tail" or inverted orientation, or in a head-to-tail orientation.

Advantageously, for example in yeast, nucleic acid encoding toxin is provided on a multicopy plasmid, such as, for yeasts 2μ (Christianson, *et al.* (1992), *Gene*, 2:110(1): 119-22), for mammalian cells a vector including oriP from Epstein Barr Virus (that may be accompanied by the initiator protein EBNA1 Kirchmaier and Sugden (1995), *J. Virol.*, 69(2): 1280-3; Wendelburg and Vos (1998), *Gene Ther.*, Oct:5(10): 1389-99), or the origin from the Bovine Papilloma Virus (that needs also two virus encoded proteins to be active (Píirsoo, *et al.* (1996), *EMBO J.*, 2:15(1): 1-11), or a viral vector.

Monocopy vectors useful in accordance with the present invention include, for yeast, ARS1 and ARSH4/CEN6 (Sikorski and Hieter (1986), *Genetics*, 122(1): 19-27; Mumberg, *et al.* (1995), *Gene*, 14:156(1): 119-22).

Suitable vectors can be chosen or constructed, containing appropriate regulatory sequences, including promoter sequences, terminator fragments, polyadenylation sequences, enhancer sequences, marker genes and other sequences as appropriate. Vectors may be plasmidic and/or viral and maintained in cells as episomes or integrated into the genome. For further details see, for example, *Molecular Cloning: a Laboratory Manual*: 2nd edition, Sambrook *et al.*, 1989, Cold Spring Harbor Laboratory Press. Many known techniques and protocols for manipulation of nucleic acid, for example in preparation of nucleic acid constructs, mutagenesis, sequencing, introduction of DNA into cells and gene

expression, and analysis of proteins, are described in detail in *Short Protocols in Molecular Biology*, Second Edition, Ausubel et al. eds., John Wiley & Sons, 1992. The disclosures of Sambrook et al. and Ausubel et al., along with
5 all other documents cited herein, are incorporated by reference.

A bacterial toxin and/or antidote or cell killing system may be provided in accordance with the present invention to a
10 eukaryotic cell selected from mammalian, human or non-human such as rabbit, guinea pig, rat, mouse or other rodent, cat, dog, pig, sheep, goat, cattle or horse, bird, such as a chicken, yeast, fungi, amphibian, fish, worm, and plant. Plants which may be employed in the present invention have
15 been noted already above.

A further aspect of the present invention provides a eukaryotic cell containing nucleic acid encoding a bacterial toxin and/or antidote or cell killing system as disclosed
20 herein, under appropriate regulatory control. The nucleic acid may be integrated into the genome (e.g. chromosome) of the cell. Integration may be promoted by inclusion of sequences which promote recombination with the genome, in accordance with standard techniques. The nucleic acid may be
25 on an extra-chromosomal vector within the cell.

A still further aspect provides a method which includes introducing the nucleic acid into a eukaryotic cell. The

introduction, which may (particularly for *in vitro* introduction) be generally referred to without limitation as "transformation", may employ any available technique. For eukaryotic cells, suitable techniques may include calcium phosphate transfection, DEAE-Dextran, electroporation, liposome-mediated transfection and transduction using retrovirus or other virus, e.g. vaccinia or, for insect cells, baculovirus.

10 The introduction may be followed by causing or allowing expression from the nucleic acid, e.g. by culturing cells (which may include cells actually transformed although more likely the cells will be descendants of the transformed cells) under conditions for expression of one or more components of the system, so that an encoded product is produced. The conditions may provide for cell killing (or inhibition of cell cycle progression, cell growth or proliferation, etc.), and/or neutralisation of the toxic effect when appropriate.

20

Introduction of nucleic acid may take place *in vivo* by way of gene therapy, as discussed below. A cell containing nucleic acid encoding a system according to the present invention, e.g. as a result of introduction of the nucleic acid into the cell or into an ancestor of the cell and/or genetic alteration of the sequence endogenous to the cell or ancestor (which introduction or alteration may take place *in vivo* or *ex vivo*), may be comprised (e.g. in the soma) within an organism which

is an animal, particularly a mammal, which may be human or non-human, yeast, fungal, amphibian, fish, worm or plant, with examples noted already above. Genetically modified or transgenic animals, birds or plants comprising such a cell are
5 also provided as further aspects of the present invention.

Thus, in various further aspects, the present invention provides a non-human animal with nucleic acid encoding a bacterial cell killing system (as disclosed) within its
10 genome. The animal may be rodent, e.g. mouse, and may provide an animal model for investigating aspects of cell cycle control, cell killing, apoptosis or other cellular process, and drug screening.

15 A further aspect provides a plant with nucleic acid encoding a bacterial cell killing system within its genome, a plant cell (which may be in culture, e.g. callus culture, or comprised in a plant or plant part), or a plant part (e.g. fruit, leaf, seed or other propagule).

20

For generation of plant material comprising nucleic acid encoding a bacterial cell killing system as disclosed, any appropriate means of transformation may be employed. Agrobacterium transformation is widely used by those skilled
25 in the art to transform both dicotyledonous and monocotyledonous species. Microprojectile bombardment, electroporation and direct DNA uptake are preferred where Agrobacterium is inefficient or ineffective. Alternatively, a

combination of different techniques may be employed, e.g. bombardment with Agrobacterium coated microparticles or microprojectile bombardment to induce wounding followed by co-cultivation with Agrobacterium. Following transformation, a
5 plant may be regenerated, e.g. from single cells, callus tissue or leaf discs, as is standard in the art. Almost any plant can be entirely regenerated from cells, tissues and organs of the plant.

10 Where a bacterial cell killing toxin is employed in accordance with the present invention there are various strategies for controlling its activity. Generally, the relevant antidote is employed to neutralise the toxic effect unless and until the toxicity is desired. Thus, for example, both toxin and
15 antidote may be expressed in normal cells, with antidote production being down-regulated in target cells (e.g. tumour cells). Toxin production may be down-regulated in normal cells and/or upregulated in target cells. Antidote production may be upregulated in normal cells and/or downregulated in
20 target cells.

Upregulation of toxin and/or antidote production, depending on context, may be achieved by a number of means. A preferred approach is to employ a promoter or other regulatory element
25 that is inducible under certain conditions, allowing for control of expression by means of application of an appropriate stimulus.

A tumour specific promoter such as telomerase RNA promoter may be employed. In plants nematode inducible promoters such as TobRB7 (Opperman et al., *Science* 263: 221-223) and PRP1 (pathogenesis related protein - see e.g. Payne et al. (1989) 5 *Plant Molecular Biology* 12: 595-596; also Memelink et al. (1990) *Plant Molecular Biology* 14: 119-126 and Payne et al. (1990) *Proc. Natl. Acad. Sci. USA* 87: 98-102) may be employed.

Downregulation of toxin and/or antidote, again depending on 10 context, may also be achieved by means of regulation of gene expression using an appropriate promoter or other regulatory element, including a repressor element, such as Tet Pr. Other approaches which may be employed include antisense regulation and ribozymes (discussed further below).

15

Thus, for example, antidote production may be downregulated by production of an antisense transcript or ribozyme. The antisense transcript or ribozyme may be produced on application of an appropriate stimulus, and may be produced 20 by expression from a sequence under transcriptional control of an inducible promoter or other regulatory element.

By "promoter" is meant a sequence of nucleotides from which transcription may be initiated of DNA operably linked 25 downstream (i.e. in the 3' direction on the sense strand of double-stranded DNA).

"Operably linked" means joined as part of the same nucleic

acid molecule, suitably positioned and oriented for transcription to be initiated from the promoter. DNA operably linked to a promoter is "under transcriptional initiation regulation" of the promoter.

5

The term "inducible" as applied to a promoter is well understood by those skilled in the art. In essence, expression under the control of an inducible promoter is "switched on" or increased in response to an applied stimulus (which may be generated within a cell or provided exogenously). The nature of the stimulus varies between promoters. Some inducible promoters cause little or undetectable levels of expression (or no expression) in the absence of the appropriate stimulus. Other inducible promoters cause detectable constitutive expression in the absence of the stimulus. Whatever the level of expression is in the absence of the stimulus, expression from any inducible promoter is increased in the presence of the correct stimulus. The preferable situation is where the level of expression increases upon application of the relevant stimulus by an amount effective to provide the desired result. Thus an inducible (or "switchable") promoter may be used which causes a basic level of expression in the absence of the stimulus which level is too low to bring about the desired result (and may in fact be zero). Upon application of the stimulus, expression is increased (or switched on) to a level which brings about the desired result.

10

15

20

25

Examples of inducible promoters for use in aspects of the present invention include a minimal promoter, such as CMV minimal promoter, fused to an enhancer for wild-type p53 activation or mutant p53 repression whether bearing the
5 consensus DNA binding sequence for wild-type p53, e.g. fragment A (Kern, *et al.* (1991), *Science*, 252(5013): 1708-11) or CON (Chen, *et al.* (1993), *Oncogene*, 8(8): 2159-66), or not, e.g. HIV 1-LTR (Subier, *et al.* (1994), *J. Virol.*, 68(1): 103-10; Gualberto and Baldwin (1995), *J. Biol. Chem.*, 25:270(34):
10 19680-3; Sawaya, *et al.* (1998), *J. Biol. Chem.*, 7:273(32): 20052-7, inducible or repressible promoters such as Tet Pr as discussed and galactose activatable GAL10-CYC1. For plants suitable promoters include the inducible GST-II promoter from maize (Jepson *et al.* (1994) *Plant Molecular Biology* 26:1855-
15 1866), alcohol inducible promoter (e.g. alc-r - see e.g. Gatz (1998) *Nature Biotechnology* 16: 140), and the Cauliflower Mosaic Virus 35S (CaMV 35S) gene promoter that is expressed at a high level in virtually all plant tissues (Benfey *et al.*, (1990) *EMBO J* 9: 1677-1684).

20

As noted, toxin production may be downregulated in non-target cells by employing elements for control of expression. Alternatively or additionally downregulation may employ antisense nucleic acid or ribozymes. Employing one or more of
25 these approaches may allow for toxin production to be eliminated in non-target cells to the extent that antidote may not be required. One or more of these approaches may be employed in addition to use of antidote to neutralise toxin

activity. Antidote may not be essential as long as appropriate control can be placed on toxin production. Where antidote is employed, antidote production itself may be controlled, as discussed.

5

In a preferred approach selectivity for expression within target cells of the toxin in accordance with the present invention is effected by a combination of (i) up-regulation of toxin production in target cells and (ii) down-regulation of
10 toxin production in non-target cells and/or neutralisation of toxin activity in non-target cells (for instance by upregulation of antidote production in non-target cells). Effect (i) will mediate the desired activity in target cells, while effect (ii) will reduce the extent of "leaky" expression
15 of that activity in non-target cells.

Where target cells are tumour cells, and non-target cells are normal cells, advantage can be taken of the fact that p53 is mutated or its function inactivated in a large proportion of
20 tumours. The p53 protein is a transcriptional activator in normal cells but is present in mutant form in a substantial proportion (40-80%) of human tumours. Even in tumours in which the p53 sequence is wild-type, its normal function in cell cycle control, DNA repair, differentiation, genome
25 plasticity or apoptosis may be abrogated, for instance by interaction with cellular protein (e.g. mdm2) or oncoviral protein (e.g. SV40 T antigen, human papillomavirus E6 protein, adenovirus E1B protein, hepatitis B virus X protein, and

Epstein-Barr BZLF-1 protein), or by being sequestered in the cytoplasm, where the p53 protein is non-functional.

Accordingly, production of the antidote (or antisense RNA or a
5 ribozyme directed against the toxin) may be controlled by a promoter whose function is upregulated by wild-type p53 in normal cells but not by mutant p53 in tumour cells. Wild-type p53 protein binds to two copies of the consensus sequence 5'-PuPuPuC(A/T)(A/T)GpyPyPy-3' (SEQ ID NO. 1) and thereby
10 transactivates the level of transcription from an operably linked promoter. Most of the mutations in the p53 gene lead to abrogation of the sequence-specific transcriptional activating function.

15 In further embodiments of the present invention, production of the toxin may be controlled by a promoter whose function is suppressed by wild-type p53 protein in normal cells, but is not suppressed or is even upregulated by mutant p53 protein, e.g. hsp70 promoter, mdm2 promoter and others. See for
20 example "The Oncogene and Tumour Suppressor Gene Facts Book", Robin Hesketh, Academic Press, Second Edition (1997) Chapter p53, pages 446-463 and references therein.

The promoters of a number of cellular genes are negatively
25 regulated by wild-type p53, include basic FGF (also activated by mutant p53), Bcl-2, human interleukin 6 and PCNA. Again, see "The Oncogene and Tumour Suppressor Gene Facts Book", Robin Hesketh, Academic Press, Second Edition (1997) Chapter

p53, pages 446-463 and references therein for examples. Viral promoters inhibited by wild-type p53 and in some cases activated by mutant versions are referenced in Deb et al. (1992) J. Virology, 66(10): 6164-6170.

5

Accordingly, such a promoter or a binding site for wild-type p53 from such a promoter may be operably linked to nucleic acid encoding the toxin. In normal cells, wild-type p53 protein suppresses production of the toxin. However, in 10 tumours where p53 is not functional and does not bind its binding site in the promoter, toxin production is derepressed.

Similarly, a response element which is activated by mutant p53 but not wildtype, such as from HIV1-LTR DNA sequences, may be 15 employed to provide for upregulation of toxin in tumour cells, or downregulation of antidote where a third component is employed to control antidote production in tumour cells. An element activated by mutant p53 element (for example) may be used to upregulate an antisense RNA, ribozyme or other factor 20 which downregulates antidote production in tumour cells.

In non-target cells production of toxin may be inhibited by using appropriate nucleic acid to influence expression by antisense regulation. Such approaches may be used to 25 downregulate antidote production in target cells. The use of anti-sense genes or partial gene sequences to down-regulate gene expression is now well-established. Double-stranded DNA is placed under the control of a promoter in a "reverse

orientation" such that transcription of the "anti-sense" strand of the DNA yields RNA which is complementary to normal mRNA transcribed from the "sense" strand of the target gene. The complementary anti-sense RNA sequence is thought then to
5 bind with mRNA to form a duplex, inhibiting translation of the endogenous mRNA from the target gene into protein. Whether or not this is the actual mode of action is still uncertain. However, it is established fact that the technique works.

10 Another possibility is that nucleic acid is used which on transcription produces a ribozyme, able to cut nucleic acid at a specific site - thus also useful in influencing gene expression. Background references for ribozymes include Kashani-Sabet and Scanlon (1995). *Cancer Gene Therapy*, 2, (3)
15 213-223, and Mercola and Cohen (1995). *Cancer Gene Therapy* 2, (1) 47-59.

Thus, an antisense RNA or ribozyme directed against toxin expression may be used to downregulate production in non-
20 target cells. Antisense RNA or ribozyme production may be placed under control of a regulatable promoter so that such production can be downregulated in target cells (for instance by means of a p53 element as discussed above).

25 An approach to downregulating toxin production in non-target cells (e.g. normal cells), and/or upregulating toxin production in target cells (e.g. tumour cells), may be instead of or in addition to regulating antidote production.

A further possibility is to use antisense RNA or a ribozyme or other approach to downregulate antidote production in target cells. Upregulating production in target cells of an antisense RNA or ribozyme against antidote may be used to
5 reduce levels of antidote in target cells and thereby increase toxin activity in those cells.

Control of translation may be employed, for instance by means of an internal ribosome entry sequence (IRES) which may be
10 controled using a RNA from yeast (Das, et al. (1996), *J. Virol.*, 70(3): 1624-32; Das, et al. (1998), *J. Virol.*, 72(7): 6638-47; Das, et al. (1998), *Front Biosci.*, 1:3: D1241-52; Venkatosan, et al. (1999), *Nucleic Acids Res.* 15:27(2): 562-72) or other that inhibit ribosome assembly at the IRES.

15

In further embodiments, the killing system, toxin and/or antidote or other inhibitor is provided to cells as protein, for instance by direct injection into target cells, such as in a tumour. In one embodiment, a carrier molecule is employed
20 to facilitate uptake by cells, e.g. a 16 aa peptide sequence derived from the homeodomain of *Antennapedia* (e.g. as sold under the name "Penetratin"), which can be coupled to a peptide via a terminal Cys residue. The "Penetratin" molecule and its properties are described in WO 91/18981. Another
25 example is VP22 (Elliott and O'Hare (1999) *Gene Ther* 6(1): 149-51; Dilber et al. (1999) *Gene Ther* 6(1): 12-21; Phelan et al. (1998) *Nat Biotechnol* 16(5): 440-3).

Expression and purification of a toxin antidote is straightforward. However, the toxic nature of a toxin such as the Kid protein makes these more difficult to over-express and purify. However, appropriate strategies are available or can
5 be devised by those of ordinary skill in the art.

Exemplifying with reference to Kis/Kid, in absence of a Kid resistant genetic background, the Kis antidote may be co-expressed at the same time in the Kid overproducer strains. The tight interaction that takes place between both proteins
10 to generate a neutralised complex allows purification from a whole bacterial extract and separation of the components afterwards by chaotropic denaturation and further chromatographic purification and renaturation of the toxic component. A bacterial one- or two- affinity chromatography-
15 based approach has been designed to purify Kid and Kid variants in high amounts and a refolding protocol has been standardised to obtain active, pure and concentrated preparations of the parD system toxin. See the experiments described below. Such an approach may be used to purify other
20 toxic components of different stability systems to be used in accordance with the present invention or other purpose.

A composition comprising nucleic acid, protein or cells according to the present invention may comprise at least one
25 additional component, such as a pharmaceutically acceptable diluent, vehicle or carrier, or a solvent or carrier for delivery to the target organism, e.g. plant.

The present invention further provides nucleic acid, proteins, cells and compositions as disclosed herein for use in a method of treatment of the human or animal body by way of therapy, e.g. for treatment of tumours, cancer, psoriasis, arteriosclerosis, any other hyper-proliferative disorder, or other disorder, the use of nucleic acid, protein, cells and compositions in the manufacture of a medicament for such treatment, and methods of treatment comprising administration of a medicament or pharmaceutical composition to a eukaryote.

Further aspects of the present invention provide methods comprising treating eukaryotic cells with nucleic acid, protein, cells or compositions as disclosed herein. The eukaryotic cells may be for example any yeast, mammalian, plant, amphibian, avian, fish or worm. Cells to be treated may be *in vitro* or in culture, or may be comprised in a mammalian (e.g. human) body or plant or plant part (e.g. fruit, leaf, seed or other propagule).

Compositions, cells and methods according to the present invention may be used in methods in which expression of a desired gene is targeted to desired cells, e.g. tumour cells as opposed to non-tumour cells. Such methods may be performed *in vivo* (e.g. by way of treatment of a human or animal body for therapeutic purposes), *ex vivo* (e.g. on cells removed from a human or animal body, prior to return of the cells to the body) or *in vitro*. Compositions and cells may be used in the manufacture of a medicament for treatment in which expression of a desired gene is targetted to target cells (e.g. tumour

cells). Nucleic acid constructs may form part of a viral vector, for instance a viral vector engineered to be suitable for administration to an individual, such as a human, and preferably additionally tumour targetting.

5

In accordance with the present invention, compositions provided may be administered to individuals. Administration is preferably in a "therapeutically effective amount", this being sufficient to show benefit to a patient. Such benefit
10 may be at least amelioration of at least one symptom. The actual amount administered, and rate and time-course of administration, will depend on the nature and severity of what is being treated. Prescription of treatment, e.g. decisions on dosage etc, is within the responsibility of general
15 practitioners and other medical doctors.

A composition may be administered alone or in combination with other treatments, either simultaneously or sequentially dependent upon the condition to be treated.

20

Pharmaceutical compositions according to the present invention, and for use in accordance with the present invention, may comprise, in addition to active ingredient, a pharmaceutically acceptable excipient, carrier, buffer,
25 stabiliser or other materials well known to those skilled in the art. Such materials should be non-toxic and should not interfere with the efficacy of the active ingredient. The precise nature of the carrier or other material will depend on

the route of administration.

Experimental support for the present invention will now be
5 described by way of illustration. Various additional aspects
and embodiments of the present invention will be apparent to
those skilled in the art.

All documents mentioned in this document are incorporated by
10 reference. "Comprising" herein is used with the meaning of
"including", that is permitting the presence of one or more
additional components or features.

EXAMPLE 1

15 *Effect of expression of the parD system in Saccharomyces
cerevisiae*

Several plasmids with different constitutive and/or
regulatable promoters were tested for their ability to express
20 both components of the parD system separately in a controlled
fashion. The results were similar with all the promoters
used. In addition to the promoters used as described in
detail in the following experiments, the inventors performed
experiments using the ADH5 promoter (constitutive; Mumberg, et
25 al. (1995), *Gene*, 14:156(1): 119-22) for kis and GAL10-CYC1
(galactose activatable Guarente, et al. (1982), *Proc. Natl.
Acad. Sci. USA*, 79(23): 7410-4) for kid.

Antidote transcription in *S. cerevisiae* was controlled by a promoter induced by Cu^{2+} , while the toxin transcription was controlled by a different promoter repressed by methionine. With that purpose, the former was cloned in a monocopy plasmid (ARSH4/CEN6 origin of replication) and the latter was cloned in a multicopy plasmid (2μ origin of replication) that confer auxotrophy for leucine and tryptophan respectively to a transfected yeast (Figure 1).

Using a multicopy plasmid for the toxin expression has two advantages: first, it reduces the possibility of selecting cells that have inactivated that protein by mutation of its DNA, as each cell should have to inactivate all the copies (10-30 molecules per haploid genome for a 2μ origin harbouring plasmid) of the kid gene present in each cell. Mutation of that gene in growth conditions in which the system is inactivated by expression of the antidote is unlikely as in that situation there is no selective pressure for the cells in order to accumulate mutations. This is verified by the fact that induction of the system exerts a clear inhibitory effect over *S. cerevisiae* growth (see below). Secondly, this approach showed that it is also possible to regulate the amount of mRNA of each component of the system by increasing or decreasing the number of encoding DNA molecules for each one (i.e. their copy number) without modifying the strength of their promoters. This allows greater flexibility in the design of systems in eukaryotes, e.g. for yeast, anti-fungals etc.

Different *S. cerevisiae* strains transfected with *kis*⁺/*kid*⁺, *kis*⁺/*kid*⁻ or *kis*⁻/*kid*⁻ plasmids were grown in liquid selective medium (-Leu/-Trp) in presence of amounts of Cu²⁺ and methionine that maintain the *parD* system in an inactivated state, before plating different serial dilutions of these cultures in solid media with a constant amount of methionine to give a constant expression of *Kid* (if any) in all the cases, but reduced concentrations of Cu²⁺ to decrease expression of its antidote from plate to plate. *Kis* and *kid* harbouring cells were not able to grow in media without Cu²⁺ and this effect is decreased as Cu²⁺ concentration increases until it reaches approximately the same rate of growth as wild type (*kis*⁻/*kid*⁻) cells. In contrast, both *kis*⁺/*kid*⁻ and wild type (*kis*⁻/*kid*⁻) cells were able to grow normally under all circumstances tested.

This experiment demonstrated that *Kid* and *Kis* are active as a toxin and its antidote respectively in yeast and that it is possible to regulate their activity (and thus *parD* activation or inactivation) by means of transcriptional control of its components in *S. cerevisiae*. It also provides indication that antidote expression alone has no side effects and that the biological process inhibited by the *parD* toxin is conserved among distantly evolved organisms.

25

EXAMPLE 2

Effect of the proteins of the parD system in Xenopus laevis

Two cell stage embryos from *Xenopus laevis* were injected at the animal pole of one of the blastomers either with Kis, Kid, both or none of them (buffer) and its effects on subsequent cell divisions were followed along time. Kid injected embryos
5 only divided correctly in the non-injected blastomer, while Kis-, Kis/Kid- and buffer- injected embryos blastomers progressed in all cases in the same way as the non-injected ones along the embryonic development stages followed in the experiment (at least until mid blastula transition, MBT).

10

This experiment further indicates that eukaryotic cell cycle progression is severely affected by non-neutralised Kid protein and suggests that this effect is not exerted in any of its gap phases (G1 or G2) as they are not present in the first
15 stages of *X. laevis* development. It also confirms that it affects a conserved biological process among distant species and offers some clues related to the possible mechanism of action of Kid (as *X. laevis* embryonic replication does not require specific DNA sequences to initiate). The fact that
20 progression through the cell cycle of the non-injected blastomers in the Kid injected embryos is not affected at all, together with the lack of effects in both halves of the other injected embryos (Kis, Kis/Kid and buffer), clearly indicates that the Kid gene product is the responsible for that
25 phenotype in eukaryotes and that the Kis gene product is responsible for its neutralisation and has no side effects *per se*, when used alone.

*EXAMPLE 3**Effect of the parD system in human cells*

The above results from yeast and amphibians show that Kid is
5 able to impede cell cycle progression through the cell cycle
in eukaryotes in a controlled fashion and that it is possible
in these organisms to substitute the prokaryotic regulatory
circuits that maintain the parD system in a silent state under
desired conditions by modulating transcription of both the
10 antidote and the toxin with different promoters.

For experiments in human cells a set of plasmids named pNATHA
(for plasmids with Neutralisable Activity that Triggers HeLa
Apoptosis) was constructed. Their mechanism of action is
15 based in the observation that in HeLa Tet Off cells a
Cytomegalovirus Early promoter (CMV Pr) maintains a constant
level of transcription of a reporter gene independently of the
presence or absence of Tetracyclin (or Doxycyclin) in the
culture medium. On the other hand, using the same cell line,
20 a Tetracyclin regulatable promoter (Tet Pr) can decrease the
level of transcription of that reporter gene by more than
three orders of magnitude upon addition of the transcriptional
regulator. In the induced state (i.e. in absence of Dox) Tet
Pr directed transcription of the reporter gene is almost two
25 orders of magnitude higher than that of the same reporter gene
under control of the CMV Pr. In the uninduced state (i.e. in
the presence of Dox), the latter transcribes almost two orders
of magnitude more efficiently than the former (Figure 2).

This transcriptional behaviour offers a window that can be used to construct the pNATHA plasmids, in which both *kis* and *kid* genes are contained in the same DNA molecule, the antidote mRNA synthesis controlled by the Tet repressible promoter and the toxin messenger levels controlled by the CMV constitutive one. Both cassettes contained Kis and Kid were cloned in either direct or inverted orientations (Figure 5). Toxin and antidote can be cloned in a tail-to-tail or tail-to-head orientation as convenient and to take advantage of transcriptional interference under appropriate control. Both may be part of the same transcriptional unit if an IRES is placed between the coding sequences.

Additional variants of both the antidote and the toxin were tested in HeLa cells, after verifying their wild type-like activity *in vivo* in *E. coli*. A Nuclear Localisation Signal (NLS) was fused to Kid and Kis to test if it would confer a more efficient effect (if any in human cells) both impeding cell cycle progression or neutralising that impedance, respectively.

All pNATHA were stably transfected in a HeLa Tet Off cell line. The *in vivo* effect of both components of the parD system on these cells was analysed before and after addition of Doxycyclin to the different cultures. The first observation of this set of experiments is that, again, after induction of the system, cell growth rate is severely inhibited in HeLa *kis*⁺/*kid*⁺ and *nlskis*⁺/*kidnls*⁺ cells. This

suggests two different things: first, that immediate transport of the toxin into the nucleus (verified by confocal microscopy of Kid immunostained samples) does not impede its toxic effect, indicating the probable nuclear localisation of its cellular target(s); and second, that the wild type components of the parD system are as active as NLS-fused ones in HeLa cells, which indicates either that entry into the nucleus is not impeded for the wild type proteins, and/or that inactivation of the cellular target(s) by Kid can occur in the cytosol. After one or two days growing in presence of Doxycyclin, and up to ten days of treatment, an induced state of parD is detectable, as kis⁺/kid⁺ cells have increased doubling time, compared to kis⁺/kid⁻ transfectants or to kis⁺/kid⁺ cells grown in absence of Doxycyclin (Figure 4). It should be noted that as only kis transcription is being modulated directly, while maintaining constant level of kid, the rate of growth for those kis⁺/kid⁺ stabilised transfectants is lower than that of their kis⁺/kid⁻ counterparts in the same conditions. This could be due to a slight escape of the system at the level of its neutralisation ability if kid transcription is not reduced selectively at the same time.

The results showed progressive reduction of cell doublings of kis⁺/kid⁺ stable transfectants upon continued exposure to Doxycyclin (i.e. to non-neutralised toxin). The inventors were interested in whether it would be possible to provide a cytostatic and/or cytotoxic effect.

Percentage of dead cells was determined after treatment with sub-lethal doses of Doxycyclin of the different stable transfectants analysed previously. As indicated before, *kis*⁺/*kid*⁻ HeLa cells showed an exponential growth rate along 5 time in both presence and absence of Doxycyclin. On the contrary, *kis*⁺/*kid*⁺ HeLa cells showed an exponential cell growth rate only when antidote transcription was maintained (i.e. in absence of Doxycyclin) but not in the opposite case, in which they reduced continuously their number of doublings 10 (Figure 5). It should be noted though that growth rate was reduced for *kis*⁺/*kid*⁻ HeLa cells grown in presence of Doxycyclin compared to that of the same stabilised cell line grown in its absence. This effect may be due to long exposure to Doxycyclin even at sub-lethal doses and, in any case, it 15 does not lead to cell death. When dead cells were counted for all the samples, *kis*⁺/*kid*⁺ HeLa cells growing in presence of Doxycyclin (i.e. in presence of non-neutralised toxin) showed a 32% and 65% of dead cells at days five and ten of treatment, respectively, while all the other samples did not show more 20 than 9% even upon ten days of treatment (Figure 5). Annexin V (i.e. an early apoptotic marker) staining of the different samples analysed, demonstrates that the observed cell death in *kis*⁺/*kid*⁺ non-neutralised HeLa cell line was due to activation of apoptosis (Figure 6).

25

MATERIALS AND METHODS

Saccharomyces cerevisiae

Plasmids

Oligonucleotides XhoIKis (5'CCGCTCGAGATGCATACCACCCGACTG3' - SEQ ID NO. 2) and KisNcoI (5'CATGCCATGGTCAGATTTCTCCTGACCAG3' - SEQ ID NO. 3) were used to amplify the kis coding region by PCR from a mini-R1 derivative. The amplified product was digested with XhoI and NcoI and cloned in the plasmid pSAL1 to construct pSAL1Kis (Mascorro-Gallardo, et al. (1996), *Gene*, 172(1): 169-70). In a similar way, oligonucleotides ATGKid (5'ATGGAAAGAGGGGAAATCTG3' - SEQ ID NO. 4) and KidEcoRI (5'CGGAATTCCTCCATGTTCAAGTC3' - SEQ ID NO. 5) were used to amplify the kid coding region using the same template and the product obtained was digested with EcoRI and cloned in the plasmid p424Met25 (Mumberg, et al. (1994), *Nucleic Acids Res.*, 25:22(25): 5767-8) digested with SmaI and EcoRI to construct the plasmid p424Met25Kid. This plasmid was amplified in a bacterial strain that overproduces Kis at the same time to abolish selection of inactivated mutants during the cloning process.

20 *In vivo assay*

Saccharomyces cerevisiae strain W303 α (MAT α , ade2-1, trp1-1, can1-100, leu2-3, 112, his3-11, ura3, psi+) was transformed with plasmids pSAL1 and p424Met25 (null), pSAL1Kis and p424Met25 (kis+/kid-) and pSAL1Kis and p424Met25kid (kis+/kid+). These cells were grown in selective medium supplemented with 500 μ M of methionine and 200 μ M of SO₄Cu to maintain the kis and kid promoters in an activated and repressed state respectively. The cultures were allowed to grow

until mid-log phase and then a 3 μ l drop of dilutions of each culture containing 15000, 1500 or 150 cells was posed in agar plates made of selective medium supplemented with 200 μ M of methionine to maintain a constant expression level of the kid gene and 0, 1, 5, 10, 20, 40, 80, 100 and 200 μ M of SO_4Cu to increase the expression level of the kis gene. The plates were incubated 48 hours at 30°C and the growth rate of each culture was analysed afterwards on each plate.

10 *Xenopus laevis*

Kis and Kid overproducers

MBPKis overproducer

Oligonucleotides ATGKis (5'ATGCATACCACCCGACTG3' - SEQ ID NO.

6) and KisEcoRI (5'TCGGAATTCAGATTTCTCCTG3' - SEQ ID NO. 7)

15 were used to amplify kis by PCR using a mini-R1 plasmid as template. The amplified product was digested with EcoRI and cloned in pMAL-c2 plasmid (Mumberg, et al. (1994), *Nucleic Acids Res.*, 25:22(25): 5767-8) between the XmnI and EcoRI sites to obtain the MBP- (Maltose Binding Protein) Kis
20 overproducer.

HisKisKid overproducer

Oligonucleotides NdeIkid (5'GGAATTCCATATGCATACCACCCGACT3' -

SEQ ID NO. 8) and kisBamHI (5'CGGGATCCTCAAGTCAGAATAGT3' - SEQ

25 ID NO. 9) were used to amplify the coding regions of kis and kid in tandem from a mini-R1 derivative. The product of PCR was digested with NdeI and BamHI and cloned in pET15b (Invitrogen) between these sites. The resultant plasmid was

digested with NcoI and BamHI and the DNA fragment codifying for Hiskiskid was purified and subcloned between these same sites in pRG-recA-NHis (Giraldo, et al. (1998), *EMBO J.*, 3:17(15): 4511-26).

5

Protein purification

MBPKis purification

Kis protein was purified as a fusion with the Maltose Binding Protein (MBP). *Escherichia coli* strain DH5 α transformed with
10 the plasmid pMBPKis was inoculated in 2 L of LB medium plus ampicillin (100 μ g/ml) at 0.04 units of Abs_{600nm} and grown with shaking at 37°C until 0.4 units of Abs_{600nm} were reached. MBPKis expression was induced then by addition of IPTG 100 μ M to the culture medium. Cells were grown for 4 hours at 37°C
15 and then pelleted in a GS3 rotor and resuspended in 10 ml lysis buffer (20 mM Tris-HCl pH 8.0, 150 mM NaCl) and frozen in liquid nitrogen. After thawing cells, 2 mg of lysozyme was added to the suspension of cells and lysis was completed by incubation at 37°C for about 10 minutes, with cooling on ice
20 every 3 minutes. A soluble fraction was obtained by addition of 40 ml of buffer 20 mM Tris-HCl pH 8.0, 600 mM NaCl and centrifugation at 30 Krpm at 4°C during 45 min in a 65 Ti rotor. MBPKis protein was purified by affinity chromatography through an amylose resin (BioLabs) following the manufacturers
25 instructions in buffer 20 mM HEPES pH 7.5, 100 mM KCl, 1 mM DTT and 10% of ethyleneglycol. MBPKis fractions were pooled and purity and concentration of the protein were determined by coomassie staining on a SDS-PAGE gel and by spectrophotometric

analysis, respectively. Fractions were stored at -80°C.

Kid purification

Escherichia coli strains C600 or TG1 transformed with the
5 overproducer pRGΔHisKisKid were grown in 2 L of LB medium plus
ampicillin (100 µg/ml) at 0.04 units of Abs_{600nm} and grown with
shaking at 37°C until 0.4 units of Abs_{600nm} were reached.
HisKis and Kid expression was induced then by addition of 25
µg/ml of nalidixic acid to the culture medium. Cells were
10 grown for 4 hours at 37°C and then pelleted in a GS3 rotor and
resuspended in 10 ml lysis buffer (20 mM Tris-HCl pH 8.0, 150
mM NaCl) and frozen in liquid nitrogen. After thawing, 2 mg
of lysozyme was added to the suspension of cells and lysis was
completed by incubation at 37°C. A soluble fraction was
15 obtained by addition of 40 ml of buffer 20 mM Tris-HCl pH 8.0,
600 mM NaCl and centrifugation at 30 Krpm at 4°C during 45 min
in a 65 Ty rotor. This soluble fraction was precipitated by
addition of 60% of ammonium sulfate and centrifugation at 40
Krpm at 4°C for 60 min. The precipitated fraction was then
20 resuspended in 1 ml of 20 mM Tris-HCl pH 7.5, 500 mM KCl) and
dialysed against the same buffer to eliminate the ammonium
sulfate. The dialysed fraction was loaded in a 5ml fast-flow
chelating sepharose (Pharmacia) activated with Ni²⁺ and
equilibrated with the dialysis buffer in which the HisKis-Kid
25 complex was retained. A gradient of 0 to 6 M of guanidinium
chloride (GnCl) in 20 mM Tris-HCl pH 7.5 was applied to the
column and denaturation of the HisKis-Kid complex bound to the
column led to retention of HisKid and elution of Kid at 5.5 M

of the chaotropic agent. Denatured Kid can be stored at -80°C until necessary. For renaturation, Kid was diluted to 5 pmol/ μ l in 6 M GnCl, 150 mM ClK, 100 mM phosphate buffer pH 6.5, 20 mM β -mercaptoethanol, 0.2 mM EDTA and 1.2 % CHAPS and 5 dialysed 5 times during 6 hours at 4°C against 200 ml (per 6 ml of protein) of 100 mM phosphate buffer pH 6.5, 150 mM KCl, 10 mM β -mercaptoethanol, 0.1 mM DTT and 10 % ethyleneglycol. The soluble and refolded protein was separated from the insoluble (denatured) one by centrifuging the mix at 40 Krpm 10 for 60 min at 4°C in a 65 Ty rotor. The supernatant was concentrated in centricon tubes (cut off 3 K) and aliquoted after determining purity and concentration of the protein by coomassie staining on a SDS-PAGE gel and spectrophotometric analysis, respectively, and stored at -80°C.

15

Embryo microinjections

MBPKis and Kid proteins were dialysed against buffer 20 mM Tri-HCl pH 8.0, 50 mM KCl and 2 μ l of MBPKid (160 ng/ μ l) and 2 μ l of MBPKis (720 ng/ μ l) were mixed with each other or with 20 μ l of dialysis buffer and incubated on ice for 10 min. 50 nl of each mix (buffer, Kis, Kid and Kis/Kid) were microinjected into dejellied two cell embryos of *Xenopus laevis* at the animal pole of one of their cells. Microinjected and non-injected embryos were then incubated in 4% of ficoll 400 in 25 MBS buffer at 18°C and allow to progress through embryonic development until stage 8-9 (blastula) was reached in the case of the non-injected controls (7-8 hours). Embryos were then photographed and the effect of microinjections analysed

afterwards.

HeLa cells

Plasmids (pNATHAs)

- 5 Oligonucleotides EcoRIKis (5'CGGAATTCATGCATACTACCACCCGACTG3' -
SEQ ID NO. 10) or EcoRINLSKis
(5'CGGAATTCATGGACAAGGTTCTAAGAAGAAGAGGAAGGTTAGCAGCATGCATACCACC
CGACTGAAG3' - SEQ ID NO. 11) and KisXbaI
(5'CTCTAGATCAGATTTCTCCTGACC3' - SEQ ID NO. 12) were used to
10 amplify kis by PCR using a mini-R1 plasmid as template. The
amplified product was digested with EcoRI and XbaI and cloned
in pTRE plasmid (Clontech) between EcoRI and XbaI sites to
obtain the pTREKis and pTRENLSKis plasmids, respectively. On
the other hand, oligonucleotides XhoIKid
15 (5'CCGCTCGAGATGGAAAGAGGGGAAATCT3' - SEQ ID NO. 13) and
KidEcoRI (SEQ ID NO. 5) were used to amplify kid by PCR using
a mini-R1 plasmid as template, and EcoRIKid
(5'CGGAATTCATGGAAAGAGGGGAAATCT3' - SEQ ID NO. 14) and
KidNLSXbaI
20 (5'GCTCTAGATCAAACCTTCCTCTTCTTCTTAGGAGGCCTGCTGCTAGTCAGAATAGTGGA
CAGGCG3' - SEQ ID NO. 15) were used with the same purpose to
obtain an NLSKid gene by PCR using a mini-R1 plasmid as
template. These two PCR products were digested with XhoI and
EcoRI or EcoRI and XbaI, respectively, and cloned between
25 these sites in the plasmid pCIneo (Promega) to obtain the
plasmids pCIneoKid and pCIneoKidNLS. These kid+ plasmids were
amplified in a bacterial strain that overproduces Kis at the
same time to abolish selection of inactivating mutants during

the cloning process. Fragment BstXI-SmaI was deleted from pClneoKid and pClneoKidNLS to eliminate the neomycin resistance gene. The resultant plasmids (pCIKid and pCIKidNLS) were digested with BgIII and BamHI and treated with 5 Klenow, and the fragment containing the kid or kidNLS genes were purified and cloned in the pTRE and pTREKis vectors digested with HindIII and treated with Klenow. For each of these constructs both orientations were selected, and plasmids pNATHA1 (kis+), pNATHA2 (kis+/kid+), pNATHA 4 (NLSkis+) and 10 pNATHA 8 (NLSkis+/kidNLS+) were obtained both in kis-kid tail-to-tail (pNATHAi) and tail-to-head (pNATHAd) orientations.

Selection of stable transfectants

5 μ g of each pNATHA was mixed with 0.5 μ g of pTKHyg plasmid 15 and HeLa Tet-Off cell line (Clontech) was transfected with these mixtures by the Lipofectamine method (Gibco). Stable transfectants were selected in DMEM medium supplemented with glutamax and 10% of tetracyclin approved fetal bovine serum (Clontech) and in the presence of 200 μ g/ml of neomycin 20 (Sigma) and 200 μ g/ml of hygromycin (Clontech) (non-toxic medium; NTM).

In vivo assays

Cell growth and death rate determination

25 HeLa Tet Off cells stably transfected with pNATHAi1+ and pNATHAi2+ were grown in NTM until they reached approximately 80% of confluency. They were trypsinised and 5×10^4 pNATHAi1+ and 2×10^4 pNATHAi2+ stably transfected cells were transferred to

4 wells of a six multiwell plate and grown for 24 hours in NTM. After that, one of the wells per sample was trypsinised and these cells pelleted and stained with trypan blue. Total and trypan blue stained (dead) cells per well were counted with a cytometer. Then, 0.1 $\mu\text{g/ml}$ of Doxycycline (Sigma) was added to the rest of wells and cells were allowed to grow in this toxic medium (TM) for 2, 5 and 10 days, changing it each 4 days when necessary but retaining the floating (dead and mitotic) cells each time that fresh TM was added.

10 Trypsinisation, trypan blue staining and counting of cells was repeated for each sample to determine the total and dead number of cells per sample.

Annexin V staining

15 HeLa Tet Off cells stably transfected with pNATHA1+ and pNATHA2+ were grown in NTM until they reached approximately 80% of confluency. They were trypsinised and 10^4 pNATHA1+ and 5×10^4 pNATHA2+ stably transfected cells were transferred to four dishes (two per sample) of 5 cm of diameter in which four polylysine coated coverslips were placed. Cells were allowed to settle down for 24 hours and then 0.1 $\mu\text{g/ml}$ of doxycyclin was added to one of the dishes per sample. Coverslips were taken out from the dishes before (day 0) and 2, 5 and 10 days after addition of doxycyclin to one of them. Fresh medium was added each 4 days if necessary. Samples growing on these coverslips were stained with FITC-Annexin V (Clontech) as suggested by the manufacturer, before fixing them, and DNA was stained with propidium iodide and Hoeschts 33258. Analysis

and counting of annexin V positive cells was done by confocal microscopy and total and apoptotic number of cells was determined.

CLAIMS

1. A method of inhibiting cell proliferation and/or cell cycle progression, the method comprising providing within
5 eukaryotic cells a bacterial toxin and an inhibitor of said toxin, optionally an antidote to the toxin wherein both toxin and antidote are proteins, under appropriate control for selective cell cycle inhibition and/or killing of target cells.
- 10 2. A method according to claim 1 wherein the cells are *in vitro*.
3. A method according to claim 1 or claim 2 wherein the
15 cells are plant cells.
4. A method according to claim 1 which is therapeutic and carried out on a human or animal body.
- 20 5. A method according to any one of the preceding claims wherein the toxin is a bacterial toxin of a post-segregational killing system.
6. A method according to any one of the preceding claims
25 wherein the toxin interferes with DNA replication, and thereby impedes cell cycle progression and/or triggers programmed cell death.

7. A method according to claim 5 or claim 6 wherein the toxin targets *DnaB*.

8. A method according to claim 7 wherein the toxin is an *E. coli* toxin.

9. A method according to claim 8 wherein the toxin is *ParD* toxin.

10 10. A method according to any one of the preceding claims wherein said toxin is provided within said cells by means of nucleic acid encoding said toxin under control of appropriate control elements for expression.

15 11. A method of inhibiting cell proliferation and/or cell cycle progression, the method comprising providing within eukaryotic cells a bacterial toxin of a post-segregational killing system.

20 12. A method according to any one of the preceding claims comprising providing to said cells said toxin and an antidote to the toxin, wherein both toxin and antidote are proteins, and controlling activity of said antidote on said toxin to control activity of said toxin on said cells.

25

13. A method according to claim 12 wherein an inhibitor of said toxin, optionally said antidote, is provided within said cells by means of nucleic acid encoding said toxin under

control of appropriate control elements for expression.

14. A method according to claim 12 or claim 13 wherein selectivity for expression said toxin within target cells is effected by a combination of (i) up-regulation of toxin production in target cells and (ii) down-regulation of toxin production in non-target cells and/or neutralisation of toxin activity in non-target cells.

15. A method according to claim 14 wherein neutralisation of toxin activity in non-target cells is effected by upregulation of antidote production in non-target cells.

16. A method according to any one of claims 12 to 15 wherein said target cells are tumour cells.

17. A method according to any one claims 11 to 16 wherein said toxin is *ParD* kid protein and said antidote is *ParD* kis protein.

20

18. A composition comprising:

(i) a bacterial toxin and an inhibitor of said toxin, optionally an antidote to said toxin wherein both toxin and antidote are proteins, or

(ii) nucleic acid encoding a bacterial toxin and an inhibitor of said toxin, optionally an antidote to said toxin wherein both toxin and antidote are proteins,

for use in a therapeutic method according to any one of

claims 4 to 17.


19. Use of a composition comprising:

(i) a bacterial toxin and an inhibitor of said toxin,
5 optionally an antidote to said toxin wherein both toxin and
antidote are proteins, or


(ii) nucleic acid encoding a bacterial toxin and an
inhibitor of said toxin, optionally an antidote to said toxin
wherein both toxin and antidote are proteins,

10 in the manufacture of a medicament for use in a
therapeutic method according to any one of claims 4 to 17.


1/6

Figure 1


2/6

Figure 2


3/6

Figure 3


4 / 6

Figure 4

5/6

Figure 5

6/6

Figure 6

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/GB 00/02743

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K38/16 A61K39/02 A61K48/00 A61K45/06 A61K31/713
A61P35/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS, CANCERLIT, CHEM ABS Data, MEDLINE, SCISEARCH

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	HAMBLETON ET AL.: "Antitoxins and botulinum toxin treatment" BRITISH MEDICAL JOURNAL, vol. 304, 1992, pages 959-960, XP000944661 * see introduction and page 960 left col.* ---	18
X	WO 94 05345 A (KABAKOV VIKTOR GRIGORIEVICH ;SELEZOV EUGENY AFANASIEVICH (RU); SKO) 17 March 1994 (1994-03-17) * see pages 26-27, 32-36 and claims 7-13 * ---	1,4,6,10
A	M. HOLCIK ET AL.: "Conditionally lethal genes associated with bacterial plasmids." MICROBIOLOGY, vol. 143, 1997, pages 3403-3416, XP000941408 * see abstract * --- -/--	1-19



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

* & * document member of the same patent family

Date of the actual completion of the international search

7 December 2000

Date of mailing of the international search report

28/12/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Gore, V

INTERNATIONAL SEARCH REPORT

Intern. Application No

PCT/GB 00/02743

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	MAGNUSON R. ET AL.: "Corepression of the P1 operon by Phd and Doc." J. BACTERIOL., vol. 180, no. 23, 1998, pages 6342-6351, XP000942967 * see abstract and page 6350 * ---	1-19
A	RAWLINGS D.E.: "Proteic toxin-antitoxin, bacterial plasmid addiction systems and their evolution with special reference to the ps system of pTF-FC2." FEMS MICROBIOL. LETTERS, vol. 176, 15 July 1999 (1999-07-15), pages 269-277, XP000942964 * see abstract * ---	1-19
X,P	WO 99 58652 A (KRISTOFFERSEN PETER ; GERDES KENN (DK); GROENLUND HUGO (DK); PEDERS) 18 November 1999 (1999-11-18) * see abstract, pages 3-4, page 8 lines 14-17, claims 51, 60, 79 and 87-88 * -----	1,2,4-6, 10-13,18

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 00/02743

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9405345 A	17-03-1994	CA 2122600 A,C	17-03-1994
		CH 685675 A	15-09-1995
		DE 4295020 C	31-07-1997
		DE 4295020 T	20-10-1994
		FR 2716940 A	08-09-1995
		JP 7500765 T	26-01-1995
		SE 511883 C	13-12-1999
		SE 9401476 A	29-04-1994
		US 5586872 A	24-12-1996
WO 9958652 A	18-11-1999	AU 3596399 A	29-11-1999